

The Millennium Galaxy Catalogue: Dynamically close pairs of galaxies and the global merger rate

Roberto De Propris

H. H. Wills Physics Laboratory, University of Bristol,
Tyndall Avenue, Bristol, BS8 1TL, United Kingdom

`R.DePropri@bristol.ac.uk`

Jochen Liske

European Southern Observatory, Karl-Schwarzschild-Straße 2,
85748 Garching b. München, Germany

`jliske@eso.org`

Simon P. Driver¹ and Paul D. Allen

Research School of Astronomy and Astrophysics, Australian National University,
Cotter Road, Weston, ACT, 2611, Australia

`spd,paul@mso.anu.edu.au`

and

Nicholas J. G. Cross

Institute for Astronomy, The University of Edinburgh, Royal Observatory, Blackford Hill,
Edinburgh, EH9 3HJ, United Kingdom

`njc@roe.ac.uk`

Received _____; accepted _____

¹PPARC Visiting Fellow, University of Bristol

ABSTRACT

We derive the number of dynamically close companions per galaxy (N_c) and their total luminosity (L_c) for galaxies in the Millennium Galaxy Catalogue: N_c is similar to the fraction of galaxies in close pairs and is directly related to the galaxy merger rate. We find $N_c = 0.0174 \pm 0.0015$ and $L_c = (252 \pm 30) \times 10^6 L_\odot$ for galaxies with $-22 < M_B - 5 \log h < -19$ with $\langle z \rangle = 0.123$ and $N_c = 0.0357 \pm 0.0027$, $L_c = (294 \pm 31) \times 10^6 L_\odot$ for galaxies with $-21 < M_B - 5 \log h < -18$, with $\langle z \rangle = 0.116$. The integrated merger rate to $z = 1$ for both samples is about 20 %, but this depends sensitively on the fraction of kinematic pairs that are truly undergoing a merger (assumed here to be 50%), the evolution of the merger rate (here as $(1+z)^3$) and the adopted timescale for mergers (0.2 and 0.5 Gyr for each sample, respectively). Galaxies involved in mergers tend to be marginally bluer than non-interacting galaxies and show an excess of both early-type and very late-type objects and a deficiency of intermediate-type spirals. This suggests that interactions and mergers partly drive the star formation and morphological evolution of galaxies.

Subject headings: galaxies: interactions

1. Introduction

In the Cold Dark Matter (CDM) framework, galaxies are assembled gradually via a process of hierarchical mergers, where increasingly more massive subunits coalesce to produce today’s luminous giant galaxies (e.g., Cole et al. 2000 and references therein): the history of mass assembly of galaxies is then reflected in the global merger rate and in its evolution with lookback time. Mergers and close interactions may also play a role in the morphological transformation of galaxies (see, for instance, Hernandez-Toledo et al. 2005; Scannapieco and Tissera 2003; Steinmetz and Navarro 2002; Junqueira et al. 1998; Mihos and Hernquist 1996), triggering of starbursts (Nikolic et al. 2004; Alonso et al. 2004; Barton et al. 2003; Bergvall et al. 2003; Lambas et al. 2003; Tissera et al. 2002; Barton et al. 2000; Donzelli and Pastoriza 1997) and fuelling of active galactic nuclei (e.g., Sanchez and González-Serrano 2003; Canalizo and Stockton 2001).

Traditionally, the observational route to measuring the merger rate has been the conventional pair fraction, under the assumption that sufficiently close galaxies will result in a merger over relatively short timescales (Zepf and Koo 1989; Burkey et al. 1994; Carlberg et al. 1994; Woods et al. 1995; Yee and Ellingson 1995; Patton et al. 1997; Wu and Keel 1998; Bundy et al. 2004); however, limited redshift information makes the derived pair fraction dependent on the mean galaxy density and the correlation function, and all such estimates are affected by contamination from unphysical pairs.

In order to address this issue, it is sometimes required that pairs also show evidence of interactions (Neuschafer et al. 1997; Le Fevre et al. 2000; Hashimoto and Oemler 2000; Xu et al. 2004): on the other hand, this introduces an element of subjectivity in the analysis, as a threshold of morphological disturbance must be chosen for objects to be considered parts of pairs. Furthermore, a number of true pairs may not show evidence of tidal tails or other deviations from symmetry (because of low surface brightness, cosmological dimming

and morphological k -corrections, for example; see discussion in Mihos 1995).

The fraction of galaxies in kinematic pairs (i.e. both spatially and dynamically close) yields a more rigorous estimate of the local pair fraction and global merger rate, although the method requires highly complete redshift information and accurate control of systematic biases arising from the flux-limited nature of redshift surveys and from incompleteness. At least for the local universe, where large redshift samples are currently available, this approach is now feasible and was first used by Patton et al. (2000, hereafter P00) who measured the number of dynamically close companions per galaxy within the range $-21 < M_B < -18$ (a statistic akin to the pair fraction) using data from the Second Southern Sky Redshift Survey (da Costa et al. 1998), and established the necessary mathematical formalism (see next section for a brief summary).

The two large redshift surveys now available, the 2dF Galaxy Redshift Survey (2dFGRS; Colless et al. 2001), and the Sloan Digital Sky Survey (SDSS; York et al. 2000), should provide a more extensive sample for such studies. Of these, the 2dFGRS is currently the largest and most complete publicly available dataset. However, because of its complicated selection function at small angular separation, which is related to restrictions on fiber placement and the survey tiling strategy, this survey is not well suited to a study of close pairs (Hawkins et al. 2003). In this paper we choose to study a smaller, but more complete dataset, which is more appropriate for the purpose of measuring the number of close companions per galaxy and the local merger rate.

The Millennium Galaxy Catalogue, hereafter MGC (Liske et al. 2003), covers a $35'$ wide and 72.2° long equatorial strip, coinciding with the northern strip of the 2dFGRS and with the SDSS Data Release 1 region (Abazajian et al. 2003), which provide $\sim 50\%$ of the MGC redshifts as well as $ugriz$ photometry. The MGC has been reimaged in the B band to deep surface brightness limits ($26 \text{ mag arcsec}^{-2}$) in order to derive accurate

structural parameters, and additional redshifts have been collected to bring the redshift completeness to 99.79% for galaxies with $B < 19.2$ and 96.05% for galaxies with $B < 20$. The total sample includes 10,095 galaxies to $B = 20$ (Driver et al. 2005). Because of its high completeness and photometric accuracy (Cross et al. 2004), the MGC is well suited to a study of the local merger rate via close pairs and to an analysis of the properties of the dynamically paired galaxies. Even the MGC, however, suffers from some incompleteness at small separations, for which we will need to make allowance in our analysis.

We describe the procedures used to derive the number of close companions per galaxy in the next section, apply these techniques to the MGC in section 3 and present the results and our discussion in section 4, dealing with theoretical interpretation of the data and with the properties of galaxies in pairs. We adopt a cosmology with $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$ and $H_0 = 100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and all absolute magnitudes used should be understood with reference to this cosmology.

2. Methodology

We follow the method of P00 and the reader is referred to that paper (and Patton et al. 2002, hereafter P02) for a more complete discussion. Here we present a summary of the technique and repeat some essential points.

Consider a primary sample of N_1 galaxies and a secondary sample of N_2 galaxies. We are interested in determining the number of galaxies from the secondary sample that are dynamically ‘close’ to a primary galaxy, per primary galaxy. This statistic, N_c , is similar to the fraction of galaxies in close pairs and is in fact identical to it for a volume-limited sample which contains only close pairs but no triplets or higher order N -tuples. We define galaxies to be dynamically close if they have a projected separation $5 < r_p < 20 \text{ kpc}$ (the

inner limit is set to avoid confusion with star formation knots or HII regions) and if they have a relative velocity of less than 500 km s^{-1} . Simulations show that such pairs would result in a merger on timescales of 10^9 years (Toomre 1977; Barnes 1988) and therefore N_c is directly related to the galaxy merger rate. These requirements have been conventionally used in the literature (e.g., Carlberg et al. 1994; Yee and Ellingson 1995; Carlberg et al. 2000) on close pair statistics.

P00 have shown that for a flux-limited sample (like the MGC) the number of close companions per galaxy is best computed as:

$$N_c = \frac{\sum_i^{N_1} w_{N_1}^i N_{c_i}}{\sum_i^{N_1} w_{N_1}^i} \quad (1)$$

and the total companion luminosity as:

$$L_c = \frac{\sum_i^{N_1} w_{L_1}^i L_{c_i}}{\sum_i^{N_1} w_{L_1}^i}. \quad (2)$$

N_{c_i} and L_{c_i} are the number and luminosity of galaxies from the secondary sample that are dynamically close to the i th primary galaxy. They are given by:

$$N_{c_i} = \sum_j w_{N_2}^j = \sum_j \frac{w_{b_2}^j w_{v_2}^j}{S_N(z_j)} \quad (3)$$

and

$$L_{c_i} = \sum_j w_{L_2}^j L_j = \sum_j \frac{w_{b_2}^j w_{v_2}^j}{S_L(z_j)} L_j, \quad (4)$$

where the sums run over those secondary galaxies that fulfill the criteria of being dynamically close to the i th primary galaxy.

The weights that are being applied to the secondary sample, w_{N_2} and w_{L_2} , have three components. The first factor corrects for the change of density of the secondary galaxies as a function of redshift due to the apparent flux limit of the sample: for a low-redshift primary galaxy we will find more close companions on average than for a higher redshift one because the companions can be drawn from a wider range of luminosities (cf. Figure 1).

The appropriate (inverse) correction factors are given by the selection functions $S_N(z)$ and $S_L(z)$ which are defined in terms of ratios of densities in flux limited vs. volume limited samples:

$$S_N(z) = \frac{\int_{M_{bright}}^{M_{lim}(z)} \Phi(M) dM}{\int_{M_{bright}}^{M_2} \Phi(M) dM} \quad (5)$$

and

$$S_L(z) = \frac{\int_{M_{bright}}^{M_{lim}(z)} \Phi(M) L(M) dM}{\int_{M_{bright}}^{M_2} \Phi(M) L(M) dM}. \quad (6)$$

$\Phi(M)$ denotes the luminosity function for which we adopt the latest MGC values from Driver et al. (2005) ($\Phi^* = 0.0177 \text{ Mpc}^{-3}$, $M^* = -19.60$ and $\alpha = -1.13$; B band). M_{lim} is the redshift dependent absolute magnitude limit:

$$M_{lim}(z) = \max(M_{faint}, m - 5 \log d_L(z) - 25 - k(z) - e(z)), \quad (7)$$

where m is the apparent magnitude limit of the survey ($B = 20$), d_L is the luminosity distance, $k(z)$ is the maximum k -correction at redshift z (P02) and $e(z)$ is of the form $-0.75 \times 2.5 \log(1 + z)$, which corresponds to a passive luminosity evolution scenario with no mergers: this form yields the same results (to within 0.01) of the $e(z) = -0.7z$ assumed by P02. Here, M_{bright} and M_{faint} are magnitude limits introduced to take account of the fact that the clustering properties of galaxies vary with luminosity (Norberg et al. 2002) and therefore we must only use galaxies in a limited luminosity range so that the clustering length does not vary too strongly.

In essence, the factors S_N and S_L in equations 3 and 4 correct the secondary sample from a flux limited sample in the range $M_{bright} < M < M_{lim}(z)$ to a hypothetical volume limited sample in the range $M_{bright} < M < M_2$.

The other two components of the w_{N_2} weights correct for boundary effects. w_{b_2} corrects for the fact that some fraction, $1 - f_b^i$, of the r_p area around the i th primary galaxy may lie outside the effective survey area: $w_{b_2}^j = 1/f_b^i$. For primary galaxies lying close to the survey

boundaries in redshift space (i.e. within 500 km s^{-1}) w_{v_2} corrects for possible companions beyond those limits: we ignore all companions between the primary galaxy and the redshift boundary and adopt $w_{v_2}^j = 2$ for those companions lying away from the boundary.

Finally we note that in equations 1 and 2 we also apply weights to the primary sample. The purpose of the w_{N_1} and w_{L_1} weights is to minimise the errors on N_c and L_c . P00 have shown that they should be chosen as:

$$w_{N_1}^i = w_{b_1}^i w_{v_1}^i S_N(z_i) \quad (8)$$

$$w_{L_1}^i = w_{b_1}^i w_{v_1}^i S_L(z_i), \quad (9)$$

where $w_{b_1}^i = f_b^i$ and $w_{v_1}^i = 0.5$ (i.e. the reciprocals of the weights for the secondary sample) for primary galaxies close to the redshift boundaries.

3. Application to the MGC

We now apply these techniques to the MGC. Figure 1 shows the absolute magnitude vs. redshift for all MGC galaxies with $B < 20$: this includes 9696 galaxies with redshifts out of 10,095, with 96.05% redshift completeness.

The mean absolute magnitude of the MGC, weighted according to equations (8) and (9) above, is $M_B = -18.5$. We then choose to analyze two subsets of the MGC data, one with $M_{bright} = -22$ and $M_2 = -19$ and another with $M_{bright} = -21$ and $M_2 = -18$ (to which ranges the results are normalized). Selection lines for both samples are drawn in Figure 1. For these samples, we use $M_{faint} = -18$ and -17 (respectively); the brighter sample contains 5756 galaxies, while the fainter one includes 6492 galaxies. The magnitude limits chosen here (as in other studies) imply that we are mostly concerned with major mergers between galaxies of approximately similar luminosity. As in P02 some galaxies are brighter than the apparent magnitude limit but blue and therefore lie ‘below’ the selection

line in Figure 1; these galaxies are excluded from the analysis to ensure that galaxies of all spectral types have an equal probability of being included within the samples.

For the bright sample ($-22 < M_B < -19$) we find a total of 137 dynamically close companions in 69 pairs¹, where two galaxies appear in two pairs. Of these companions, 60 lie in the magnitude range $-19 < M_B < -18$ (i.e. between M_2 and M_{faint} for this sample. For the faint sample ($-21 < M_B < -18$) we find 176 companions in 89 pairs, with one triple and where two galaxies appear in two pairs each. Of these, 27 lie between M_2 and M_{faint} . This is similar to P00 in that galaxies between these two limits do not contribute significantly to the pair fraction for the $\sim L^*$ galaxies; however, these objects are much more significant contributors for the brighter sample. This may be evidence that the brighter galaxies tend to have fainter companions.

Applying equations 1 and 2 we find $N_c = 0.0147 \pm 0.0013$ and $L_c = 213 \pm 25 \times 10^6 L_\odot$ for galaxies with $-22 < M_B < -19$, with mean $z = 0.124$ and $N_c = 0.0301 \pm 0.0023$, with $L_c = (248 \pm 26) \times 10^6 L_\odot$ for galaxies with $-21 < M_B < -18$ with mean $z = 0.116$, where the errors are derived by jackknife resampling of the data (Efron 1982).

Figure 2 shows postage stamps for a random selection of pairs; about half or more of these galaxies show clear signs of interaction such as disturbed morphologies, rings and tidal features, demonstrating that our sample indeed includes *bona fide* physical pairs. The full list of companions provides a useful objectively selected sample of galaxies in the early phase of a merger. Table 1 lists all unique pairs found in MGC; in order the columns show: the MGC ID of the primary galaxy, the IDs of the other pair members, the RA and Dec. (2000) for the primary galaxy and its radial velocity.

¹One of the pairs contributes only a single companion because r_p is just below 20 kpc at the redshift of one of the galaxies but just above 20 kpc at the redshift of the other

However, we know that the MGC has some redshift incompleteness. We proceed to estimate the number of missed pairs in the following way. We repeat our analysis including galaxies without redshifts and using only the r_p criterion, but requiring that if a galaxy without redshift is involved in a pair, its absolute magnitude (assigning it the redshift of the other object) be consistent with the absolute magnitude limits being used (i.e. it must lie between the selection lines shown in Figure 1). This yields an extra 36 companions for the $-22 < M_B < -19$ sample and 47 companions for the $-21 < M_B < -18$ sample. Since some of these pairs may be due to chance superposition, we repeat our analysis using only galaxies with redshifts, but solely using the r_p criterion. We find a total of 193 and 262 companions, respectively, of which we know 137 and 176 to be ‘real’. The fraction of ‘true’ companions is therefore 70% and we apply this to derive an extra 25.2 companions for the first sample and 32.9 companions for the second sample. This yields a correction of 18.4% and 18.7% to the derived values of N_c and L_c (for the brighter and fainter sample, respectively) owing to galaxies whose redshifts are missing from the survey.

These values are larger than one might expect from the low redshift incompleteness ($\sim 4\%$) of the survey; the main source of incompleteness lies in galaxies that were never targeted and these objects lie preferentially at close angular separations to other galaxies: since the MGC derives most of its redshifts from 2dF observations (including 2dFGRS data) and SDSS surveys, the corrections are affected by the instrumental bias against close pairs due to fiber placement constraints (although this is lower than in the 2dFGRS because of the higher MGC completeness).

4. Discussion

After applying the incompleteness correction derived above we find that $N_c = 0.0174 \pm 0.0015$ and $L_c = (252 \pm 30) \times 10^6 L_\odot$ for galaxies with $-22 < M_B < -19$ and

$N_c = 0.0357 \pm 0.0027$ and $L_c = (294 \pm 31) \times 10^6 L_\odot$ for galaxies with $-21 < M_B < -18$, with mean redshifts as given above. If we assume that the merger rate evolves as $(1+z)^m$ with $m \sim 3$ (Le Fevre et al. 2000; Gottlöber et al. 2001; Patton et al. 2002) we find that our N_c of 0.0357 ± 0.0027 translates to 0.0257 ± 0.0019 at the mean SSRS2 redshift of 0.015, which is in good agreement with the number of companions of 0.0225 ± 0.0052 measured by P00 for galaxies with $-21 < M_B < -18$. At $z = 0.289$, P02 find $N_c = 0.0334 \pm 0.0081$ for these same galaxies, converted to our cosmology (D. Patton, private communication), which corresponds to $N_c = 0.0217 \pm 0.053$ at our mean $z = 0.116$.

4.1. Comparison with theoretical models

Predictions of the expected merger rate in CDM models can be derived from numerical simulations. However, these simulations are usually conducted at much lower resolution than the typical pair separation used here. Although it is possible to extrapolate to higher resolution, by resampling the larger simulations, this procedure is suspicious as information may already have been lost on small scales (Gottlöber et al. 2001). Additionally, it is difficult to relate dark haloes to their luminous content: it is generally assumed that each galaxy corresponds to a single halo, but this assumption is likely to be too simplistic.

Khochfar and Burkert (2001) calculate the fraction of close pairs and the evolution of the merger rate in a Λ CDM cosmology. The merger fraction they derive for galaxies of $M \approx 3 \times 10^{12} M_\odot$ (which corresponds to $-21 < M_B < -18$) is approximately in agreement with our result and P00, but their models cannot fully reproduce the drop in the merger fraction by a factor of 2 for the brighter galaxies. The simple ‘chance hypothesis’, where galaxies fall into each other’s gravitational influence zone simply by random motions, appears to work somewhat better: galaxies with $-22 < M_B < -19$ have an approximately 6 times lower chance to be in close proximity than galaxies with $-21 < M_B < -18$, by

integration of the luminosity function, but such objects are also ~ 3 times more clustered (Norberg et al. 2002). Therefore the pair fraction should be lower by about a factor of 2 which is in reasonable agreement with our findings. This is apparently in contrast with the conclusions of Xu et al. (2004) who appear to reject the chance hypothesis for dwarfs in the proximity of giants; the hypothesis may only apply to giant galaxies, which are more resilient than dwarfs. Clearly, a more consistent interpretation of these results must await a more thorough theoretical modelling of this phenomenon.

Since most companions are found in pairs, rather than in triplets or higher order N -tuples, our derived value of N_c is comparable to the fraction of galaxies in close pairs. Following Yee and Ellingson (1995) and P00 we now assume that half of our *dynamically* close pairs are actually *physically* close pairs and will hence merge. We now integrate the merger rate to $z = 1$ to derive the fraction of present day galaxies that have undergone a major merger since this time. The fraction of merger remnants is (P00):

$$f_{rem} = 1 - \prod_{k=1}^N \frac{1 - F_{mg}(z_k)}{1 - 0.5F_{mg}(z_k)} \quad (10)$$

where $F_{mg}(z)$ is the merger rate at redshift z , which is assumed to vary as $F_{mg}(z) = F_{mg}(0)(1+z)^m$, with $m = 3$, z_k corresponds to a lookback time of $t = kT_{mg}$ and T_{mg} is the merger timescale. Following P00, we assume that this merger timescale is 0.5 Gyr for galaxies with $-21 < M_B < -18$ and, scaling by the luminosity of the galaxies, 0.2 Gyr for galaxies with $-22 < M_B < -19$. We therefore have 27 merger timescales to $z \sim 1$ for the brighter sample and 11 timescales for the fainter sample. This implies that $\sim 22.7\%$ of $-22 < M_B < -19$ galaxies and $\sim 19.2\%$ of $-21 < M_B < -18$ galaxies have undergone a major merger since $z \sim 1$, approximately the last half of the Hubble time. P00 derive a remnant fraction $\sim 6.6\%$ for their sample, but they assume a slightly different cosmology and $m = 0$, while Lin et al. (2004) also obtain a remnant fraction of $\sim 10\%$, but use

$m = 0.51$ and P02 derived an integrated merger fraction of 15% with $m = 2.3$. Note, however, that these values are strongly dependent on the assumed merger timescales, index for the evolution of the merger rate and fraction of merging galaxies. For the sake of comparison, Λ CDM models predict that about 50% of $L > L^*$ galaxies have undergone a major merger since $z \sim 1$ (Murali et al. 2002), which is about a factor of 2 lower than our estimate.

4.2. The properties of galaxies in pairs

It is interesting to compare the properties of galaxies in pairs with respect to their parent sample. Since only $\sim 20\%$ of these objects have undergone a merger in the last half of the Hubble time, this should provide a sample of ‘undisturbed’ galaxies and offer some insight as to the effects of close interactions (and mergers) on galaxy morphology and star formation. Figure 2 already offers a hint of this: many galaxies show obvious signs of interactions, with tidal tails, rings and other signs of morphological disturbance, a connection exploited by the asymmetry parameter (Conselice et al. 2003).

Figure 3 shows the colour distribution in rest $u - r$ for galaxies in both samples, compared with that of their parent distribution as well as the distribution of morphologies in the pairs and parent samples. Although the errors on the distributions for paired galaxies are large, because of small number statistics, there is some evidence that galaxies in pairs tend to be bluer than those in the parent sample, by approximately 0.1 mag in $u - r$, suggesting that close interactions induce star formation episodes (Alonso et al. 2004; Nikolic et al. 2004; Barton et al. 2003; Lambas et al. 2003; Barton et al. 2000), although the average blueing for our sample is more consistent with relatively mild starbursts (Bergvall et al. 2003). On the other hand, not all of our kinematic pairs may be real merging systems and therefore, not all objects may show enhanced star formation.

Comparison of the morphologies (determined from visual examination of the images) shows that there are marginally more E/S0 galaxies and Sd/Irr galaxies among pairs, while there are fewer Sabc galaxies. The excess of early-type galaxies in pairs was noted by Junqueira et al. (1998), and this was attributed to an excess of lenticulars by Rampazzo and Sulentic (1992) and Rampazzo et al. (1995) who suggest that pairs containing early-type members may be the result of early merging of groups (which would have additional companions handy to fuel further merging). The excess of very late-type galaxies may be explained by interactions increasing the star formation rate and therefore leading to later classifications. These may be long-lived interacting systems rather than objects in an early phase of a merger (Junqueira et al. 1998).

Mergers and interactions therefore alter the morphologies of the member galaxies and induce star formation episodes. Scenarios in which local conditions (i.e. mergers and close encounters) are primarily responsible for morphological evolution and the decline of star formation in denser environments have been recently proposed following analysis of large local samples of field and cluster galaxies (Balogh et al. 2004; De Propris et al. 2004; Croton et al. 2005), although the change in the fraction of blue galaxies is much higher than the merger fraction we derive (suggesting a large contribution from lower luminosity galaxies). Wider and deeper surveys would be valuable, in allowing consideration of the importance of minor mergers.

RDP is supported by a PPARC fellowship. JL acknowledges an ESO fellowship. The Millennium Galaxy Catalogue consists of imaging data from the Isaac Newton Telescope and spectroscopic data from the Anglo Australian Telescope, the ANU 2.3m, the ESO New Technology Telescope, the Telescopio Nazionale Galileo, and the Gemini Telescope. The survey has been supported through grants from the Particle Physics and Astronomy Research Council (UK) and the Australian Research Council (AUS). The data and data

products are publicly available from <http://www.eso.org/~jliske/mgc/> or on request from JL or SPD. We would like to thank the referee, D. R. Patton, for a number of helpful comments and suggestions that made the paper clearer and better.

REFERENCES

- Abazajian, K. et al. 2003, AJ, 126, 2081
- Alonso, M. S., Tissera, P. B., Coldwell, G. and Lambas, D. G. 2004, MNRAS, 352, 1081
- Balogh, M. L. et al. 2004, MNRAS, 348, 1355
- Barton, E., Geller, M. J. and Kenyon, S. J. 2000, ApJ, 530, 660
- Barton, E., Geller, M. J. and Kenyon, S. J. 2003, ApJ, 582, 668
- Barnes, J. E. 1988, ApJ, 331, 699
- Bergvall, N., Laurikainen, E. and Aalto, S. 2003, A&A, 405, 31
- Bundy, K. A., Fukugita, M., Ellis, R. S., Kodama, T. and Conselice, C. J. 2004, ApJ, 601, L123
- Burkey, J. M., Keel, W. C., Windhorst, R. A. and Franklin, B. E. 1994, ApJ, 429, L13
- Canalizo, G. and Stockton, A. 2001, ApJ, 555, 719
- Carlberg, R. G., Pritchet, C. J. and Infante, L. 1994, ApJ, 435, 40
- Carlberg, R. G. et al. 2000, ApJ, 532, L1
- Cole, S., Lacey, C. G., Baugh, C. M. and Frenk, C. S. 2000, MNRAS, 319, 168
- Colless, M. M. et al. 2001, MNRAS, 328, 1039
- Conselice, C., Bershadsky, M. A., Dickinson, M. and Papovich, C. 2003, AJ, 126, 1183
- Cross, N. G. J., Driver, S. P., Liske, J., Lemon, D. J., Peacock, J. A., Cole, S., Norberg, P. and Sutherland, W. J. 2004, MNRAS, 349, 576

- Croton, D. J. et al. 2005, MNRAS, 356, 1155
- da Costa, L. N. et al. 1998, AJ, 116, 1
- De Propris, R. et al. 2004, MNRAS, 351, 125
- Donzelli, C. J. and Pastoriza, M. G. 1997, ApJS, 111, 181
- Driver, S. P., Liske, J., Cross, N. G. C., De Propris, R. and Allen, P. D. 2005, MNRAS, 360, 81
- Efron, B. 1982, The Jackknife, the Bootstrap and other resampling plans, CMBS-NSF Regional Conference Series in Applied Mathematics (Philadelphia: Society for Industrial and Applied Mathematics)
- Gottlöber, S., Klypin, A., Kravtsov, A. V. 2001, ApJ, 546, 223
- Hashimoto, Y. and Oemler, A. 2000, ApJ, 530, 652
- Hawkins, E. et al. 2003, MNRAS, 346, 78
- Hernandez-Toledo, H. M., Avila-Reese, V., Conselice, C. J. and Puerari I. 2005, AJ, 129, 682
- Junqueira, S., de Mello, D. F. and Infante, L. 1998, A&AS, 129, 69
- Khochfar, S. and Burkert, A. 2001, ApJ, 561, 517
- Lambas, D. G., Tissera, P. B., Alonso M. S. and Coldwell, G. 2003, MNRAS, 346, 1189
- Le Fevre, O. et al. 2000, MNRAS, 311, 565
- Lin, L. et al. 2004, ApJ, 617, L9

- Liske, J., Lemon, D. J., Driver, S. P., Cross, N. J. G. and Couch, W. J. 2003, MNRAS, 344, 307
- Mihos, J. C. 1995, ApJ, 438, L75
- Mihos, J. C. and Hernquist, L. 1996, ApJ, 464, 641
- Murali, C., Katz, N., Hernquist, L., Weinberg, D. and Davé, R. 2002, ApJ, 571, 1
- Neuschafer, L. W., Im, M., Ratnatunga, K. U., Griffiths, R. E. and Casertano, S. 1997, ApJ, 480, 59
- Nikolic, B., Cullen, H. and Alexander P. 2004, MNRAS, 355, 874
- Norberg, P. et al. 2002, MNRAS, 332, 827
- Patton, D. R., Pritchet, C. J., Yee, H. K. C., Ellingson, E. and Carlberg, R. G. 1997, ApJ, 475, 29
- Patton, D. R., Carlberg, R. G., Marzke, R. O., Pritchet, C. J., da Costa, L. N. and Pellegrini, P. S. 2000, ApJ, 536, 153 (P00)
- Patton, D. R. et al. 2002, ApJ, 565, 208 (P02)
- Rampazzo, R. and Sulentic, J. W. 1992, A&A, 259, 43
- Rampazzo, R., Reduzzi, L., Sulentic, J. W. and Madejsky, R. 1995, A&AS, 110, 131
- Sanchez, S. F. and González-Serrano, J. I. 2003, A&A, 406, 435
- Scannapieco, C. and Tissera, P. P. 2003, MNRAS, 338, 880
- Steinmetz, M. and Navarro, J. F. 2002, New Astronomy, 7, 155

- Tissera, P. B., Domnguez-Tenreiro, R., Scannapieco, E. and Sáiz, A. 2002, MNRAS, 333, 327
- Toomre, A. 1977, in The Evolution of Galaxies and Stellar Populations, ed. B. M. Tinsley and R. B. Larson (New Haven: Yale University Press), p. 401
- Woods, D., Fahlman, G. G. and Richer, H. B. 1995, ApJ, 454, 32
- Wu, W. and Keel, W. C. 1998, AJ, 116, 1513
- Xu, C. K., Sun, Y. C. and He, X. T. 2004, ApJ, 603, L73
- Yee, H. K. C. and Ellingson, E. 1995, ApJ, 445, 37
- York, D. et al. 2000, AJ, 120, 1579
- Zepf, S. E. and Koo, D. C. 1989, ApJ, 337, 34




Fig. 1.— The absolute magnitude-redshift distribution of MGC galaxies, showing the selection lines for the $-22 < M_B < -19$ and the $-21 < M_B < -18$ samples in the upper and lower panel, respectively. In each panel, the thick solid line delimits the magnitude limited sample with $M_{\text{bright}} < M_B < M_{\text{lim}}(z)$ used in the analysis, while the dashed box shows the hypothetical volume limited sample with $M_{\text{bright}} < M_B < M_2$ to which the results are




Fig. 2.— Postage stamps of a random selection of pair galaxies. The stamps are $40h^{-1}$ kpc on the side and are centered on a randomly chosen pair member.




Fig. 3.— Histograms of the colour and morphology distributions for galaxies in pairs and for their parent samples.

Table 1. MGC: Unique close pairs and triples

MGC ID	Comp. ID	RA (2000)	Dec. (2000)	<i>cz</i>
533	537	10:00:23.9	00:00:51.9	18918
919	921	10:01:17.9	-00:11:02.9	27348
956	960,963	10:01:28.3	-00:12:44.1	18525
973	978	10:01:08.3	-00:13:32.7	28017
977	95002	10:01:02.9	-00:13:38.8	27411
1579	1580	10:03:44.2	00:01:50.5	19611
2858	2866	10:08:56.4	-00:02:16.5	28737
3118	3121	10:09:03.8	-00:16:42.8	20574
3526	3530	10:11:21.4	-00:10:07.3	20907
4397	4742	10:15:53.3	00:05:07.4	22011
4420	4421	10:16:28.6	-00:01:34.1	31554
4450	4452	10:15:44.1	-00:03:02.7	42861
4484	4486	10:16:23.3	-00:04:49.7	28458
4651	4655	10:16:25.1	-00:14:25.2	21162
4988	4995	10:17:48.6	-00:01:50.3	18621
5974	5977	10:22:07.5	-00:03:39.6	21354
6551	6552	10:22:58.6	-00:10:23.7	33912
6610	6614	10:23:35.5	-00:13:17.3	28611
6888	6895	10:26:13.8	-00:02:00.8	31461
6892	6895	10:26:13.9	-00:02:04.3	31029
7876	7881	10:28:56.3	-00:02:02.0	28410
8169	8170	10:30:11.0	00:00:51.2	33249

Table 1—Continued

MGC ID	Comp. ID	RA (2000)	Dec. (2000)	<i>cz</i>
12189	12190	10:49:20.1	00:14:17.8	33327
13718	13720	10:54:50.9	-00:09:16.4	32703
15486	15490	11:03:31.6	-00:10:27.1	22404
16242	16245	11:07:38.2	-00:01:59.2	19728
16245	16250	11:07:38.2	-00:02:13.1	19776
16408	16413	11:07:44.9	-00:11:21.5	20094
17921	18001	11:15:22.0	00:06:55.2	8193
18822	18824	11:17:50.8	00:03:07.6	29955
19025	19318	11:19:51.1	00:05:12.4	31302
19565	19579	11:20:57.1	-00:05:03.7	7506
19604	19605	11:20:53.2	-00:07:29.6	29520
21115	21118	11:29:46.1	00:14:33.1	31263
22241	22246	11:32:40.1	00:01:35.7	39351
22591	22594	11:35:55.1	-00:16:07.4	19590
22594	22595	11:35:56.6	-00:16:12.0	19698
23246	23247	11:40:15.9	-00:09:59.2	22626
25070	25072	11:49:47.9	00:07:42.7	23514
25500	25502	11:51:36.3	00:00:02.2	18150
25601	25604	11:52:26.6	-00:06:30.2	38721
25660	25662	11:50:51.4	-00:10:53.7	31017
25897	25900	11:54:31.3	00:10:20.2	32496
26507	96408	11:56:19.5	-00:12:18.3	32637

Table 1—Continued

MGC ID	Comp. ID	RA (2000)	Dec. (2000)	<i>cz</i>
26526	26527	11:56:01.7	-00:13:04.7	32535
26724	26725	11:57:37.6	00:07:32.0	32235
26970	26973	11:56:37.8	-00:11:10.0	31950
27383	27386	12:00:28.9	-00:07:24.5	24288
27476	27477	11:58:41.6	-00:12:47.6	28482
27627	27629	12:01:46.0	00:14:32.2	31470
27897	27898	12:00:44.7	-00:09:22.2	49578
29266	29269	12:06:43.0	-00:12:13.7	27825
29405	29407	12:09:41.2	00:12:06.2	29826
29901	29904	12:12:09.9	00:10:29.7	37899
31392	31394	12:16:58.5	-00:13:18.4	21576
31991	95166	12:21:56.5	00:09:48.7	31875
32019	32022	12:21:48.2	-00:00:23.4	32475
32110	32111	12:22:17.9	-00:07:43.1	51849
33182	33189	12:27:22.7	00:06:43.2	47955
35114	35124	12:35:42.4	-00:12:53.9	6873
37672	37676	12:47:42.5	-00:08:14.3	26871
37916	37918	12:50:28.1	00:13:36.3	13989
37988	37990	12:50:31.9	00:09:30.3	24855
39047	39048	12:53:02.0	-00:16:12.6	25461
39373	39375	12:54:37.8	-00:11:04.4	24822
40301	40302	13:02:21.9	00:14:40.3	20403

Table 1—Continued

MGC ID	Comp. ID	RA (2000)	Dec. (2000)	<i>cz</i>
41961	41965	13:07:02.4	00:03:24.2	23976
44577	44581	13:18:41.2	-00:10:25.8	14487
45563	45565	13:22:47.1	-00:13:47.4	24699
45798	45804	13:26:34.4	00:09:09.8	25878
46553	45565	13:27:49.0	-00:15:34.1	42756
49445	49450	13:40:17.2	00:02:06.0	43479
52736	52742	13:53:26.2	-00:09:04.2	31308
52767	52768	13:53:18.7	-00:10:16.1	56688
53371	53376	13:55:41.2	-00:14:44.5	31623
53554	53560	13:58:41.9	00:14:51.5	9942
53797	95689	13:56:46.2	-00:08:54.1	32229
56147	56149	14:05:02.9	-00:15:05.0	9777
59005	59009	14:16:17.2	-00:09:25.2	25059
59318	59324	14:17:59.6	00:16:02.8	15963
59445	59446	14:16:45.1	-00:01:14.3	37581
59700	59721	14:17:37.2	00:03:50.6	15882
60716	60719	14:20:56.7	-00:04:23.6	30654
60869	60904	14:22:27.5	00:03:38.5	9729
61615	61618	14:24:10.7	00:01:47.2	37986
64206	64334	14:34:21.9	00:06:27.2	41031
64512	64513	14:33:10.6	-00:08:07.2	40839
64512	64515	14:33:10.6	-00:08:07.2	40839

Table 1—Continued

MGC ID	Comp. ID	RA (2000)	Dec. (2000)	<i>cz</i>
68038	68048	14:44:04.5	00:03:14.6	22059
95060	95061	12:08:11.7	00:03:50.2	29916
96246	96247	14:27:30.7	-00:10:38.6	48996
96944	96945	14:15:24.6	00:16:02.2	53166